

**#0055**

Effect of diet on insulin resistance in adults at risk of developing type 2 diabetes mellitus

**Author/s:**

Yethindra Vityala<sup>1</sup>; Rithwik Goud Burri<sup>2</sup>; Krishna Chaitanya Meduri<sup>2</sup>; Sai Praneeth Duwuri<sup>2</sup>; Sriharsha Lankala<sup>2</sup>; Annie Monica Ch<sup>2</sup>; Raghu Ram Shanmukh Nemani<sup>3</sup>; Tejaswi Vadde<sup>4</sup>; Sai Prasad Karuturi<sup>4</sup>; Harish Kumar Jella<sup>4</sup>

**Organizations/Affiliations:**

<sup>1</sup>AJ Institute of Medical Sciences and Research Centre Mangalore, Karnataka, India; <sup>2</sup>Maheshwara Medical College and Hospital, Hyderabad, Telangana, India; <sup>3</sup>GSL Medical College and General Hospital, Rajahmundry, Andhra Pradesh, India; <sup>4</sup>SVS Medical College, Mahbubnagar, Telangana, India

**Abstract**

Introduction: Insulin resistance (IR) has been associated with poor dietary practices and is considered to be the basis of the pathophysiology of type 2 diabetes mellitus (T2DM).

Objective: To examine the association between diet and IR in adults at risk of developing T2DM.

Methods: A cross-sectional study of 112 individuals at high risk of developing T2DM (mean age: 43.4 years; 59 [52.7%] women) was done between May and September 2023. BMI and HOMA-IR were calculated using the participants' weight, height, glucose, and insulin levels, respectively. Using the Health Diet Score to assess diet and the HOMA-IR to calculate IR, the IR cutoff value of 2.5 was determined. To investigate the association between diet quality and IR, the Pearson correlation coefficient and linear regression were used.

Results: HOMA-IR is negatively correlated with diet quality (OR -0.42; 95% CI (-0.56, -0.22);  $p < 0.001$ ), in model 1, adjusted for age (OR -0.44; 95% CI (-0.73, -0.25);  $p = 0.003$ ), in model 2, adjusted for age and BMI (OR -0.32; 95% CI (-0.61, 0.03);  $p = 0.042$ ), and in model 3, adjusted for age, BMI, and gender (OR -0.37; 95% CI (-0.54, 0.12);  $p = 0.068$ ). Diet quality is significantly explained by 0.6% by the HOMA-IR, considering age ( $R^2 = 0.006$ ,  $p = 0.003$ ), where there is no significance and only adjustment is added for gender ( $R^2 = 0.083$ ,  $p = 0.076$ ).

**Conclusion:** Individuals at risk of developing T2DM had a negative correlation between diet quality and IR. Considering adult age, gender, and BMI, the HOMA-IR explains 6.1% of the variation in diet quality.

**Keywords:**

Insulin resistance; type 2 diabetes mellitus; diet; diet quality.

**Abbreviations:**

Insulin resistance (IR); type 2 diabetes mellitus (T2DM).

**Funding and Conflicts of interest:**

No financial support was received for the study and the authors declare no conflicts of interest.